

Neutron Star Properties from an NJL Model Modified to Simulate Confinement

S. Lawley^a W. Bentz^b and A. W. Thomas^c

^aSpecial Research Centre for the Subatomic Structure of Matter,
University of Adelaide, Adelaide SA 5005, Australia

^bDepartment of Physics, School of Science, Tokai University, 117 Kita-Kaname, Hiratsuka 259-1207,
Japan

^cJefferson Lab, 12000 Jefferson Avenue, Newport News, VA 23606, U.S.A.

The NJL model has recently been extended with a method to simulate confinement. This leads in mean field approximation to a natural mechanism for the saturation of nuclear matter. We use the model to investigate the equation of state of asymmetric nuclear matter and then use it to compute the properties of neutron stars.

1. Introduction

The properties of isospin asymmetric matter at finite density are of particular interest in the study of nuclear physics and neutron stars. In particular, there is great interest in exploring whether stars with sufficiently high core density contain some form of quark matter [1,2,3]. In order to investigate such questions it is natural to start with a model for dense matter which is built from the quark level. A number of models of this kind have been constructed, for example the quark-meson coupling (QMC) model [4,5,6,7,8] and the chiral soliton model [9].

We work within the framework of the flavor SU(2) Nambu–Jona-Lasinio (NJL) model, modified to simulate quark confinement. As explained by Bentz and Thomas [10], a description of nuclear matter can be obtained, which naturally exhibits the property of saturation, thus allowing for stable nuclei at low densities. The nucleon is constructed as a quark-diquark state by solving the Faddeev equation in the static approximation [11]. Using the proper time cut-off scheme to regularize the integrals [12,13], effectively introduces the phenomena of confinement, while leading to the stabilization of symmetric nuclear matter [10]. The general form for the equa-

tion of state is derived from the quark-level Lagrangian using the path integral formalism [14]. This model has been developed further to incorporate the phase transition to quark matter, including a color superconducting state [14]. However, in the present work we will deal with central densities where such transitions have not yet occurred.

In this work we consider isospin asymmetric matter (neutrons, protons and electrons) in β -equilibrium. Electrons are introduced to the system as a Fermi gas, balancing the positive charge of the protons. We calculate neutron star masses, radii and profiles by solving the Tolman–Oppenheimer–Volkoff (TOV) equations [15]. We find that this model can support stable stars up to a maximum mass of $2.19M_{\odot}$ where the central density is $0.97fm^{-3}$. A typical mass neutron star ($1.4M_{\odot}$) has a central density of $0.4fm^{-3}$, and a radius around 12.4 km.

2. Model for the Equation Of State

The Lagrangian for the flavor SU(2) NJL model has the general form

$$\mathcal{L} = \bar{\psi}(i \not{\partial} - m)\psi + \sum_{\alpha} G_{\alpha}(\bar{\psi}\Gamma_{\alpha}\psi)^2, \quad (1)$$




Figure 1. Equation of state for asymmetric nuclear matter for various values of the proton fractions (Y_p).




Figure 2. Proton fraction for charge-neutral nuclear matter (neutrons, protons and electrons) in chemical equilibrium.

where m is the current quark mass, ψ is the flavor SU(2) quark field, Γ_α are the matrices in Dirac, flavor and color space, characterizing the chirally symmetric 4-fermi interaction between the quarks, and G_α are the coupling constants associated with these interactions. More explicitly, in matter of density ρ , we have

$$\begin{aligned} \mathcal{L} = & \bar{\psi}(i\partial - M - 2G_\omega\gamma^\mu\omega_\mu - G_\rho\gamma^\mu\tau\cdot\rho_\mu)\psi - \\ & \frac{(M-m)^2}{4G_\pi} + G_\omega\omega^\mu\omega_\mu + G_\rho\rho^\mu\rho_\mu + \mathcal{L}_I, \end{aligned} \quad (2)$$

where we define the effective quark mass (i.e., the constituent quark mass in-medium) as $M = m - 2G_\pi\langle\rho|\bar{\psi}\psi|\rho\rangle$ and the vector and isovector fields in the nuclear medium are given by $\omega^\mu = \langle\rho|\bar{\psi}\gamma^\mu\psi|\rho\rangle$ and $\rho^\mu = \langle\rho|\bar{\psi}\gamma^\mu\tau\psi|\rho\rangle$. From this Lagrangian we derive both the equation of state and the quark-diquark structure of the bound “nucleon”. The interaction Lagrangian $\mathcal{L}_I$ consists of a sum of qq channel interaction terms, which are isolated using Fierz transformations [16]

$$\mathcal{L}_I = \mathcal{L}_{I_s} + \mathcal{L}_{I_a} + \dots \quad (3)$$

In order to construct the nucleon state, we restrict ourselves to the scalar diquark channel ($\mathcal{L}_{I_s}$) in

the present calculation. This will soon be extended to include vector di-quark correlations. The equation of state can be derived using the path integral formalism [14], or equivalent Lagrangian techniques [17]. The energy density is expressed as

$$\mathcal{E} = \mathcal{E}_{vac} + \mathcal{E}_N + \mathcal{E}_\omega + \mathcal{E}_\rho, \quad (4)$$

where the vacuum term is given by

$$\begin{aligned} \mathcal{E}_{vac} = & 12i \int \frac{d^4k}{(2\pi)^4} \ln \frac{k^2 - M^2 + i\epsilon}{k^2 - M_0^2 + i\epsilon} + \frac{(M-m)^2}{4G_\pi} \\ & - \frac{(M_0-m)^2}{4G_\pi}, \end{aligned} \quad (5)$$

The Fermi motion of the nucleons contributes

$$\mathcal{E}_N = \sum_N \gamma_N \int \frac{d^3k}{(2\pi)^3} \theta(k_{F_N} - k) \sqrt{k^2 + M_N^2}, \quad (6)$$

where M_N is the effective mass of the bound nucleon. (This is calculated by solving the Faddeev equations using the in-medium masses of both the constituent quark and scalar di-quark.) The sum here is over neutrons and protons ($N = n, p$), which give equal contributions in the case of symmetric nuclear matter (of Fermi momentum k_{F_N}).

For nuclear matter at rest the omega meson con-




Figure 3. Equations of state for neutron matter (solid line), matter in β - equilibrium (dashed line) and symmetric nuclear matter (dotted line).

tribution is given by,

$$\mathcal{E}_\omega = 9G_\omega \rho_B^2, \quad (7)$$

where the baryon density, ρ_B , is the sum of the proton and neutron densities, ρ_p and ρ_n , respectively and G_ω is the vector coupling constant. The contribution to the energy density from the rho meson is given by,

$$\mathcal{E}_\rho = 9G_\rho (\rho_p - \rho_n)^2, \quad (8)$$

where the coupling constant G_ρ is calculated at the empirical saturation density, $(\rho_B, E_B/A) = (0.16 \text{ fm}^{-3}, -15 \text{ MeV})$, so that the symmetry energy coefficient is 32.5 MeV .

For a given density the constituent quark mass M in Eq. (5) is determined by the condition $\partial \mathcal{E} / \partial M = 0$, where the quark mass at zero density is $M_0 = 400 \text{ MeV}$. The nucleon mass M_N in Eq. (6) is a function of M , determined by the pole in the Faddeev equation [10]. The energy density and pressure are calculated implicitly as a function of baryon density. As shown in




Figure 4. Neutron star masses as a function of central baryon density.

Fig. 1, the presence of protons softens the equation of state. This effect leads to more compact neutron stars. To find the equation of state for matter in β -equilibrium, we introduce electrons, $\mathcal{E}_e = k_{F_e}^4 / 4\pi^2$, and impose the conditions of charge neutrality ($\rho_p = \rho_e$) and chemical equilibrium ($\mu_e = \mu_n - \mu_p$). Fig. 2 indicates the proton fraction for matter in β -equilibrium as a function of baryon density.

Table 1

Parameters for the calculations of the properties of the nucleon and nuclear matter.

$G_\pi [GeV^{-2}]$	19.60
$G_s [GeV^{-2}]$	$0.51 G_\pi$
$G_\omega [GeV^{-2}]$	$0.37 G_\pi$
$G_\rho [GeV^{-2}]$	$0.092 G_\pi$
$\Lambda_{UV} [MeV]$	638.5
$\Lambda_{IR} [MeV]$	200
$m [MeV]$	16.93




Figure 5. Neutron star mass versus radius for pure neutron matter and for nuclear matter in β -equilibrium.

3. Results and Discussion

Given the relationship between $\mathcal{E}$ and P we can integrate the TOV equations,

$$\frac{dP}{dr} = -\frac{G(\mathcal{E}(r) + P(r))(4\pi r^3 P(r) + M(r))}{r(r - 2GM(r))} \quad (9)$$

$$M = 4\pi \int_0^R \mathcal{E}(r) r^2 dr \quad (10)$$

The stars generated from matter in β -equilibrium are similar to those composed of pure neutron matter. Fig. 6 illustrates the profiles of typical mass stars for each equation of state. In the case of matter in β -equilibrium the central density is a little less than 3 times normal nuclear matter density. This corresponds to a proton fraction of about 15% (see Fig. 2), in the centre of the star.

4. Concluding Remarks

In conclusion, we have seen that matter in β -equilibrium produces more compact stars than




Figure 6. Pressure profiles for neutron stars of mass $1.4 M_\odot$.

pure neutron matter. The abundance of protons depends on the value of G_ρ , which in our case is relatively low. While the differences are quite small, the effect is clear. We have found that this model predicts densities of about $0.4 fm^{-3}$ in the core of typical mass neutron stars. Results for masses and radii are in basic agreement with astrophysical data.

This work was supported by the Australian Research Council and DOE contract DE-AC05-84ER40150, under which SURA operates Jefferson Lab.

REFERENCES

1. N. K. Glendenning and F. Weber, *Astrophys. J.* **559** (2001) L119 [arXiv:astro-ph/0003426].
2. F. Weber, arXiv:astro-ph/0407155.
3. P. K. Panda, D. P. Menezes and C. Providencia, *Phys. Rev. C* **69** (2004) 025207 [arXiv:nucl-th/0310075].
4. P. A. M. Guichon, *Phys. Lett. B* **200** (1988) 235.
5. P. A. M. Guichon, K. Saito, E. N. Rodionov and A. W. Thomas, *Nucl. Phys. A* **601** (1996) 349 [arXiv:nucl-th/9509034].
6. K. Saito and A. W. Thomas, *Phys. Rev. C* **51** (1995) 2757 [arXiv:nucl-th/9410031].

7. P. K. Panda, G. Krein, D. P. Menezes and C. Providencia, Phys. Rev. C **68** (2003) 015201 [arXiv:nucl-th/0306045].
8. G. Chanfray, Nucl. Phys. A **721** (2003) 76 [arXiv:nucl-th/0212085].
9. J. R. Smith and G. A. Miller, arXiv:nucl-th/0407093.
10. W. Bentz, A. W. Thomas, Nucl. Phys. A 696 (2001) 138.
11. N. Ishii, W. Bentz and K. Yazaki, Phys. Lett. B **318** (1993) 26.
12. G. Hellstern, R. Alkofer and H. Reinhardt, Nucl. Phys. A **625** (1997) 697 [arXiv:hep-ph/9706551].
13. D. Ebert, T. Feldmann and H. Reinhardt, Phys. Lett. B **388** (1996) 154 [arXiv:hep-ph/9608223].
14. Nucl. Phys. A **720** (2003) 95 [arXiv:nucl-th/0210067].
15. J. R. Oppenheimer, G. M. Volkoff, Phys. Rev. 55 (1939) 374.
16. N. Ishii, W. Bentz, K. Yazaki, Nucl. Phys. A **587** (1995) 617.
17. B. D. Serot, J. D. Walecka, in *Advances in Nuclear Physics*, ed by J. W. Negele and E. Vogt (Plenum Press, New York, 1986), Vol. 16, 1.
18. N. Fröhlich, H. Baier, W. Bentz, Phys. Rev. C **57** (1998) 3447.
19. U. Vogl, W. Weise, Prog. Part. Nucl. Phys. **27** (1991) 195.
20. N. K. Glendenning, *Compact Stars*, (Springer, New York, 2000).